

Periodic weak solutions for a classical one-dimensional isotropic biquadratic Heisenberg spin chain

Boling Guo^a, Ming Zeng^{b,*}, Fengqiu Su^c

^a Center for Nonlinear Studies, Institute of Applied Physics and Computational Mathematics, PO Box 8009-28, Beijing 100088, China

^b College of Applied Sciences, Beijing University of Technology, PingLe Yuan 100, Chaoyang District, Beijing 100022, China

^c Department of Mathematical and Computer Science, Center University of Nationality, Beijing 100081, China

Received 22 March 2005

Available online 12 September 2006

Submitted by M.D. Gunzburger

Abstract

In this paper, we consider the model proposed by Lakshmanan, Porsezian and Daniel [M. Lakshmanan, K. Porsezian, M. Daniel, Effect of discreteness on the continuum limit of the Heisenberg spin chain, *Phys. Lett. A* 133 (1988) 483–488] describing the continuum isotropic biquadratic Heisenberg spin chain in one dimensional case. Following the results of integrability in [K. Porsezian, M. Daniel, M. Lakshmanan, On the integrability aspects of the one-dimensional classical continuum isotropic Heisenberg spin chain, *J. Math. Phys.* 33 (5) (1992) 1807–1816], we prove the existence of periodic weak solutions to this model.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Isotropic biquadratic Heisenberg spin chain; Weak solutions; Viscosity vanishing method

1. Introduction

In 1988, Lakshmanan, Porsezian and Daniel [1] gave the following Hamiltonian for a classical one-dimensional isotropic biquadratic Heisenberg spin chain with N spins characterized by nearest neighbor bilinear and biquadratic exchange interaction:

$$H = -J \sum_{ij} [S_i \cdot S_j + k(S_i \cdot S_j)^2], \quad i, j = 1, 2, \dots, N. \quad (1.1)$$

* Corresponding author.

E-mail address: mengnan218@sohu.com (M. Zeng).

Here the spins S_i 's are three-dimensional vectors; J and k denote the bilinear and biquadratic exchange parameters, respectively. The equation of motion corresponding to (1.1) with nearest neighbor interacting spins is

$$\frac{dS_n}{dt} = JS_n \times \{(S_{n+1} + S_{n-1}) + k[S_{n+1}(S_n \cdot S_{n+1}) + S_{n-1}(S_n \cdot S_{n-1})]\}. \quad (1.2)$$

In order to consider the continuum limit of (1.2) in the long wavelength and low temperature, the authors allow the lattice parameter a between nearest neighbor sites to approach zero and assume a slow variation of S_n over a lattice distance. By introducing $u(x, t) = u(na, t) = S_n(t)$, u can be expanded in a Taylor series up to fourth-order. After rescaling the time variable appropriately, (1.2) takes the form

$$u_t = u \times \left\{ u_{xx} + \gamma_1 u_{xxx} + \gamma_2 \left[(u \cdot u_{xx}) u_{xx} + \frac{2}{3} (u \cdot u_{xxx}) u_x \right] \right\}, \quad (1.3)$$

where $\gamma_1 = a^2/12$ and $\gamma_2 = ka^2/(1+2k)$. When $\gamma_1 = \gamma_2 = 0$, (1.3) turns into the well-known Landau–Lifshitz equation [3]. When $\gamma_1 = 0$ and $\gamma_2 \leq 0$, the corresponding equation has been deduced by Fizev [4] when he revisited the one-dimensional classical compressible Heisenberg chain, which was also considered previously by Cieplak and Turski [5]. Furthermore, Guo et al. [6] proved the existence of measure-valued solutions. Since the fourth-order term $S \times S_{xxxx}$ on the right side of (1.3) seems to be dissipative from the mathematical point of view, we want to derive a solution stronger than the one considered in [6], that is just the main result of this paper.

However, the method we will use here cannot be applied for all γ_1 and γ_2 , in fact, it is related to the integrability of (1.3). For this purpose, Porsezian et al. [2] turned (1.3) into its equivalent form through a differential geometric approach. It reads

$$iq_t + q_{xx} + 2|q|^2 q + \gamma_1 q_{xxx} - 4\delta_1 |q|^2 q_{xx} - 4\delta_2 |q|^2 q_{xx}^* - 4\delta_3 q |q_x|^2 - 4\delta_4 q^* |q_x|^2 - 24\delta_5 |q|^4 q = 0, \quad (1.4)$$

where $\delta_1 = 3\gamma_1 + 2\gamma_2$, $\delta_2 = 2\gamma_1 + \gamma_2$, $\delta_3 = 9\gamma_1 + 4\gamma_2$, $\delta_4 = \frac{7}{2}\gamma_1 + 2\gamma_2$ and $\delta_5 = 3\gamma_1 + \gamma_2/2$. (1.4) can be viewed as a generalization of the nonlinear Schrödinger equation because (1.4) reduces to the nonlinear Schrödinger equation when $\gamma_1 = \gamma_2 = 0$.

Porsezian et al. [2] utilized Painlevé singularity structure analysis to conclude that (1.4) is integrable if and only if $\gamma_2 = -\frac{5}{2}\gamma_1$. Consequently, (1.3) takes the form

$$u_t = u \times \left\{ u_{xx} + \gamma_1 u_{xxx} - \frac{5}{2}\gamma_1 \left[(u \cdot u_{xx}) u_{xx} + \frac{2}{3} (u \cdot u_{xxx}) u_x \right] \right\}, \quad (1.5)$$

Daniel et al. [7] obtained the similar result on the nonlinear spin dynamics of an anisotropic Heisenberg ferromagnetic spin chain with octupole-dipole interaction.

Without loss of generality, we will assume γ_1 equals to 1.

By virtue of

$$\begin{aligned} |u|^2 = 1 &\Rightarrow u \cdot u_x = 0 \\ &\Rightarrow u \cdot u_{xx} = -|u_x|^2 \\ &\Rightarrow u \cdot u_{xxx} = -\frac{3}{2}(|u_x|^2)_x, \end{aligned}$$

we can rewrite (1.5) as

$$u_t = u \times u_{xxx} + \left[\left(1 + \frac{5}{2}|u_x|^2 \right) u \times u_x \right]_x. \quad (1.6)$$

We will consider the following periodic initial-value problem:

$$\begin{cases} u_t = u \times u_{xxx} + \left[\left(1 + \frac{5}{2} |u_x|^2 \right) u \times u_x \right]_x, & \text{in } \Omega \times (0, T), \\ u(x, 0) = u_0(x), & \text{in } \Omega, \\ u(x + 2D, t) = u(x, t), & \text{in } \mathbb{R} \times [0, T), \end{cases} \quad (1.7)$$

where $\Omega = (-D, D)$ and D is a positive constant.

By the viscosity vanishing method, we can derive the following theorem.

Theorem 1.1. *Let $u_0 \in H^2(\Omega)$, $|u_0| = 1$ a.e. Then there exists a weak solution for (1.7) such that*

- (i) $u \in L^\infty(0, T; H^2(\Omega))$, $|u| = 1$ a.e.;
- (ii) for any $\varphi \in C^\infty(\bar{\Omega} \times [0, T])$, $\varphi(x, T) = 0$, $\varphi(-D, t) = \varphi(D, t)$, there holds

$$\begin{aligned} - \int_{\Omega \times (0, T)} u \cdot \varphi_t dx dt &= \int_{\Omega} u_0 \varphi(x, 0) dx + \int_{\Omega \times (0, T)} (u \times u_{xx}) \cdot \varphi_{xx} dx dt \\ &\quad + 2 \int_{\Omega \times (0, T)} (u_x \times u_{xx}) \cdot \varphi_x dx dt \\ &\quad + \int_{\Omega \times (0, T)} \left(1 + \frac{5}{2} |u_x|^2 \right) u \times u_{xx} \cdot \varphi dx dt \\ &\quad + \int_{\Omega \times (0, T)} 5(u_x \cdot u_{xx}) u \times u_x \cdot \varphi dx dt, \end{aligned}$$

where T depends on $\|u_0\|_{H^2}$.

The paper is organized as follows: In Section 2 we consider (1.7) with an additional ε viscosity term, see (2.1); In Section 3 we first give a uniform estimate on the solution u_ε of (2.1) with respect to ε , then we take the limit $\varepsilon \rightarrow 0$ to obtain the weak solution of (1.7).

2. An ε -parameter equation

Consider the following equation:

$$\begin{cases} u_t = u \times u_{xxx} + \left[\left(1 + \frac{5}{2} |u_x|^2 \right) u \times u_x \right]_x - \varepsilon u_{xxxx} \\ \quad - \varepsilon \left(\frac{3}{2} |u_x|_{xx}^2 + u_x \cdot u_{xxx} \right) u, \\ u(x, 0) = u_0, \\ u(x + 2D, t) = u(x, t). \end{cases} \quad (2.1)$$

We have the following lemma about smooth solutions for (2.1).

Lemma 2.1. *If u is a smooth solution of (2.1), then $|u| = 1$.*

Proof. Let us set $v(x, t) = |u(x, t)|^2$, then v satisfies

$$\begin{cases} v_t = -\varepsilon v_{xxx} - \varepsilon \left(\frac{3}{2} |u_x|_{xx}^2 + u_x \cdot u_{xxx} \right) (v - 1), \\ v(x, 0) = 1, \\ v(x + 2D, t) = v(x, t). \end{cases}$$

Furthermore we set $w = v - 1$, then w satisfies

$$w_t = -\varepsilon w_{xxx} - G(x, t)w, \quad (2.2)$$

$$w(x, 0) = 0, \quad (2.3)$$

$$w(x + 2D, t) = w(x, t), \quad (2.4)$$

where $G(x, t) = \frac{3}{2} |u_x|_{xx}^2 + u_x \cdot u_{xxx}$. Taking the scalar product with w in (2.2), and integrating over Ω , we obtain

$$\frac{1}{2} \frac{d}{dt} \int_{-D}^D |w|^2 dx + \varepsilon \int_{-D}^D |w_{xx}|^2 dx \leq \varepsilon \max_{x,t} |G(x, t)| \int_{-D}^D |w|^2 dx.$$

From Gronwall's inequality and $w(x, 0) = 0$, we deduce that $w(x, t) = 0$, that is, $|u(x, t)| = 1$. \square

Next we use the difference method to prove the existence of solutions for (2.1).

To proceed, we need the following classical notations:

$$u_h = \{u_j = u(x_j) \mid j = 0, 1, \dots, J\}, \quad x_j = jh, \quad h = 2D/J,$$

$$\|u_h\|_{L^\infty} = \sup_{j \in \{0, 1, \dots, J\}} |u_j|,$$

$$\|u_h\|_p = \left(\sum_{j=0}^{J-1} |u_j|^p h \right)^{1/p},$$

$$\|\delta^k u_h\|_p = \left(\sum_{j=0}^{J-k} \left| \frac{\Delta_+^k u_j}{h^k} \right|^p h \right)^{\frac{1}{p}},$$

where $2 \leq p < \infty$.

With these notations, we have the following lemmas.

Lemma 2.2. [8] *Let q, r be real numbers and j, m be integers such that $1 \leq q, r \leq \infty$, $0 \leq j < m$. If $u \in W^{m,r}(\Omega) \cap L^q(\Omega)$, then*

$$\|D^j u\|_p \leq C \|u\|_q^{1-\alpha} \|u\|_{m,r}^\alpha,$$

where $\|\cdot\|_p = \|\cdot\|_{L^p(\Omega)}$, $\|\cdot\|_{m,r} = \|\cdot\|_{W^{m,r}(\Omega)}$, $p \geq 1$, $\frac{j}{m} \leq \alpha \leq 1$ and

$$\frac{1}{p} - j = \frac{1-\alpha}{q} + \alpha \left(\frac{1}{r} - m \right).$$

Lemma 2.3. [9] Let p be a real number and j, m be integers such that $2 \leq p \leq \infty$, $0 \leq j < m$. Then

$$\|\delta^j u_h\|_p \leq C \|u\|_2^{1-\alpha} \left(\|\delta^m u_h\|_2 + \frac{\|u_h\|_2}{(2D)^m} \right)^\alpha,$$

where $\alpha = \frac{1}{m}(j + \frac{1}{2} - \frac{1}{p})$.

Lemma 2.4. [9] Let $u_h = \{u_j \mid j = 0, \pm 1, \dots, \pm J, \dots\}$, $v_h = \{v_j \mid j = 0, \pm 1, \dots, \pm J, \dots\}$ such that

$$u_{j+J} = u_j,$$

$$v_{j+J} = v_j.$$

Then

- (i) $\sum_{j=0}^{J-1} u_j \Delta_+ v_j = -\sum_{j=1}^J v_j \Delta_- u_j$;
- (ii) $\sum_{j=1}^J u_j \Delta_+ \Delta_- v_j = -\sum_{j=0}^{J-1} \Delta_+ u_j \Delta_+ v_j$;
- (iii) $\Delta_+(u_j v_j) = u_{j+1} \Delta_+ v_j + v_j \Delta_+ u_j$,

where Δ_+ , Δ_- denote the forward and backward difference, respectively.

Now we can establish the following difference-differential equation corresponding to (2.1) ($\varepsilon = 1$):

$$\begin{aligned} \frac{du_j}{dt} = & u_j \times \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} + \left(1 + \frac{5}{2} \left| \frac{\Delta_+ u_j}{h} \right|^2 \right) u_j \times \frac{\Delta_+ \Delta_- u_j}{h^2} \\ & + 5 \left(\frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+ \Delta_- u_j}{h^2} \right) u_j \times \frac{\Delta_+ u_j}{h} - \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} \\ & - \left(3 \left| \frac{\Delta_+ \Delta_- u_j}{h^2} \right|^2 + 4 \frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+^2 \Delta_- u_j}{h^3} \right) u_j, \end{aligned} \quad (2.5)$$

with

$$u_j|_{t=0} = u_{0j} = u_0(jh), \quad (2.6)$$

$$u_{j+J} = u_j, \quad (2.7)$$

where $j = 0, \pm 1, \dots, \pm J, \dots$, $h = 2D/J$, $J > 0$.

It is obvious that the ordinary differential equation (2.5) admits a local smooth solution. For this solution, we shall give some estimates uniformly in h , then pass to the limit $h \rightarrow 0$ to get a local smooth solution of (2.2).

Lemma 2.5. If $u_j(x) \in H^2(\Omega)$, then there exist constants $T > 0$ and $C > 0$ independent of h such that

$$\sup_{0 \leq t \leq T} \|u_h\|_2 \leq C,$$

$$\sup_{0 \leq t \leq T} \|\delta u_h\|_2 \leq C,$$

$$\sup_{0 \leq t \leq T} \|\delta^2 u_h\|_2 \leq C.$$

Proof. (1) Taking the scalar product with $u_j h$ in (2.5) and summing up over j from 1 to J lead to

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \sum_{j=1}^J |u_j|^2 h &= - \sum_{j=1}^J u_j \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} \\ &\quad - \sum_{j=1}^J \left(3 \left| \frac{\Delta_+ \Delta_- u_j}{h^2} \right|^2 + 4 \frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+^2 \Delta_- u_j}{h^3} \right) |u_j|^2 h. \end{aligned} \quad (2.8)$$

Applying Lemmas 2.3, 2.4 and Young's inequality, we get

$$\frac{d}{dt} \sum_{j=1}^J |u_j|^2 h + \|\delta^2 u_h\|_2^2 \leq \frac{1}{4} \|\delta^3 u_h\|_2^2 + C(1 + \|u_h\|_2^{12} + \|\delta u_h\|_2^4 + \|\delta^2 u_h\|_2^4). \quad (2.9)$$

(2) Taking the scalar product with $(\Delta_+ \Delta_- u_j)/h$ in (2.5) and summing up over j from 1 to J lead to

$$\begin{aligned} \sum_{j=1}^J u_{jt} \frac{\Delta_+ \Delta_- u_j}{h} &= \sum_{j=1}^J u_j \times \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} \cdot \frac{\Delta_+ \Delta_- u_j}{h^2} h \\ &\quad + 5 \sum_{j=1}^J \left(\frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+ \Delta_- u_j}{h^2} \right) u_j \times \frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+ \Delta_- u_j}{h^2} h \\ &\quad - \sum_{j=1}^J \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} \cdot \frac{\Delta_+ \Delta_- u_j}{h^2} h \\ &\quad - \sum_{j=1}^J \left(3 \left| \frac{\Delta_+ \Delta_- u_j}{h^2} \right|^2 + 4 \frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+^2 \Delta_- u_j}{h^3} \right) u_j \cdot \frac{\Delta_+ \Delta_- u_j}{h^2} h. \end{aligned}$$

Applying Lemmas 2.3, 2.4, Young's inequality and Hölder's inequality, we get

$$\frac{d}{dt} \|\delta u_h\|_2^2 + \frac{3}{2} \|\delta^3 u_h\|_2^2 \leq C(1 + \|u_h\|_2^{12} + \|\delta u_h\|_2^8 + \|\delta^2 u_h\|_2^{10}). \quad (2.10)$$

(3) Taking the scalar product with $(\Delta_+^2 \Delta_-^2 u_j)/h^3$ in (2.5) and summing up over j from 1 to J lead to

$$\begin{aligned} \sum_{j=1}^J u_{jt} \frac{\Delta_+^2 \Delta_-^2 u_j}{h^3} &= \sum_{j=1}^J \left(1 + \frac{5}{2} \left| \frac{\Delta_+ u_j}{h} \right|^2 \right) u_j \times \frac{\Delta_+ \Delta_- u_j}{h^2} \cdot \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} h \\ &\quad + \sum_{j=1}^J 5 \left(\frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+ \Delta_- u_j}{h^2} \right) u_j \times \frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} h \\ &\quad - \sum_{j=1}^J \left| \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} \right|^2 h \\ &\quad - \sum_{j=1}^J \left(3 \left| \frac{\Delta_+ \Delta_- u_j}{h^2} \right|^2 + 4 \frac{\Delta_+ u_j}{h} \cdot \frac{\Delta_+^2 \Delta_- u_j}{h^3} \right) u_j \cdot \frac{\Delta_+^2 \Delta_-^2 u_j}{h^4} h. \end{aligned}$$

Applying Lemmas 2.3, 2.4, Young's inequality and Hölder's inequality, we get

$$\frac{d}{dt} \|\delta^2 u_h\|_2^2 + \frac{3}{4} \|\delta^3 u_h\|_2^2 \leq C(1 + \|u_h\|_2^{32} + \|\delta u_h\|_2^{40} + \|\delta^2 u_h\|_2^{16}). \quad (2.11)$$

Adding (2.9)–(2.11) up, after a simple calculation we have

$$\|\delta^k u_h\|_2 \leq C, \quad \int_0^T \|\delta^{k+2} u_h\|_2^2 \leq C, \quad k = 0, 1, 2, \quad 0 \leq t \leq T,$$

where T depends on D , $\|u_{0h}\|_2$, $\|\delta u_{0h}\|_2$ and $\|\delta^2 u_{0h}\|_2$.

By induction and a limiting procedure, we can eventually derive a local smooth solution on $[0, T]$ for (2.1). \square

3. Uniform estimates and the limiting case

We have indeed proved the existence of smooth solutions u_ε for the ε -equation (2.1). In addition, we need the uniform estimates on u_ε with respect of ε . To demonstrate the method used in this section, we begin with the estimate on the solution for the original equation (1.7).

Lemma 3.1. *If u is a smooth solution for (1.7), then for any $T_1 > 0$, there exists a constant $C > 0$ depending only on $\|u_0\|_{H^2}$ such that*

$$\sup_{0 \leq t \leq T_1} \|u\|_{H^2(\Omega)} \leq C.$$

Proof. (1) Taking the scalar product with u in (1.7), we have

$$\frac{d}{dt} |u|^2 = 0,$$

which implies that $|u| \equiv |u_0| = 1$.

(2) Taking the scalar product with $-u_{xx}$ in (1.7) and integrating over Ω , we have

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|u_x\|_2^2 &= \int_{\Omega} u \times u_{xx} \cdot u_{xxx} dx - 5 \int_{\Omega} (u_x \cdot u_{xx})(u \times u_x \cdot u_{xx}) dx = I_1 - I_2, \\ I_1 &= - \int_{\Omega} (u_x \times u_{xx}) \cdot u_{xxx} dx. \end{aligned} \quad (3.1)$$

Since $|u| = 1$, we have $u \cdot u_x = 0$. Thus u , u_x and $u \times u_x$ consist of an orthonormal basis of R^3 . u_{xx} can be formulated as

$$u_{xx} = \alpha_1 u + \alpha_2 u_x + \alpha_3 u \times u_x, \quad (3.2)$$

where $\alpha_1 = u \cdot u_{xx} = -|u_x|^2$, $\alpha_2 = (u_x \cdot u_{xx})/|u_x|^2$ and $\alpha_3 = (u \times u_x \cdot u_{xx})/|u_x|^2$.

Substituting $\alpha_1, \alpha_2, \alpha_3$ into (3.2), we have

$$u_x \times u_{xx} = |u_x|^2 u \times u_x + (u \times u_x \cdot u_{xx}) u. \quad (3.3)$$

So

$$I_1 = - \int_{\Omega} (u \times u_x \cdot u_{xxx}) |u_x|^2 dx - \int_{\Omega} (u \times u_x \cdot u_{xx})(u \cdot u_{xxx}) dx. \quad (3.4)$$

If we differentiate $u \cdot u_x = 0$ twice, we have

$$u \cdot u_{xxx} = -\frac{3}{2}|u_x|_x^2. \quad (3.5)$$

For the first term on the right-hand side of (3.4), we have

$$-\int_{\Omega} (u \times u_x \cdot u_{xxx})|u_x|^2 dx = -\int_{\Omega} (u \times u_x \cdot u_{xx})_x |u_x|^2 dx = -\int_{\Omega} (u \times u_x \cdot u_{xx})|u_x|_x^2 dx. \quad (3.6)$$

Substituting (3.5) and (3.6) into (3.4), we have

$$I_1 = 5 \int_{\Omega} (u \times u_x \cdot u_{xx})(u_x \cdot u_{xx}) dx = I_2,$$

which means

$$\frac{d}{dt} \|u_x\|_2^2 = 0 \Rightarrow \|u_x(t)\|_2 = \|u_x(0)\|_2, \quad 0 < t < T_1. \quad (3.7)$$

(3) Taking the scalar product with u_{xxx} in (1.7) and integrating on Ω , we have

$$\frac{1}{2} \frac{d}{dt} \|u_{xx}\|_2^2 = \int_{\Omega} \left(1 + \frac{5}{2}|u_x|^2\right) u \times u_{xx} \cdot u_{xxx} dx + 5 \int_{\Omega} (u_x \cdot u_{xx})(u \times u_x \cdot u_{xxx}) dx. \quad (3.8)$$

We also have

$$\begin{aligned} \frac{1}{4} \frac{d}{dt} \|u_x\|_4^4 &= \int_{\Omega} |u_x|^2 u_x \cdot u_{xt} dx \\ &= -\int_{\Omega} |u_x|^2 u_{xx} \cdot u_t dx - 2 \int_{\Omega} (u_x \cdot u_{xx})(u_x \cdot u_t) dx \\ &= -\int_{\Omega} |u_x|^2 u_{xx} \cdot (u \times u_{xxx} + 5(u_x \cdot u_{xx})u \times u_x) dx \\ &\quad - 2 \int_{\Omega} (u_x \cdot u_{xx})u_x \cdot \left(u \times u_{xxx} + \left(1 + \frac{5}{2}|u_x|^2\right)u \times u_{xx}\right) dx \\ &= -\int_{\Omega} |u_x|^2 u_{xx} \cdot (u \times u_{xxx}) dx - 2 \int_{\Omega} (u_x \cdot u_{xx})u_x \cdot (u \times u_{xxx}) dx \\ &\quad - 2 \int_{\Omega} (u_x \cdot u_{xx})u_x \cdot (u \times u_{xx}) dx \\ &= \int_{\Omega} |u_x|^2 u \times u_{xx} \cdot u_{xxx} dx + 2 \int_{\Omega} (u_x \cdot u_{xx})u \times u_x \cdot u_{xxx} dx \\ &\quad - 2 \int_{\Omega} (u_x \cdot u_{xx})u_x \cdot (u \times u_{xx}) dx. \end{aligned} \quad (3.9)$$

The last term on the right-hand side of (3.9) is equal to $\frac{2}{5} \int_{\Omega} u \times u_{xx} \cdot u_{xxx} dx$ from step (2), so comparing (3.9) with (3.8), we have

$$4 \frac{d}{dt} \|u_{xx}\|_2^2 = 5 \frac{d}{dt} \|u_x\|_4^4. \quad (3.10)$$

Integrating (3.10) from 0 to T_1 , we have

$$4 \|u_{xx}(T_1)\|_2^2 - 5 \|u_x(T_1)\|_4^4 = \|u_{0xx}\|_2^2 - 5 \|u_{0x}\|_4^4. \quad (3.11)$$

Applying Lemma 2.2, we have

$$\|u_x\|_4^4 \leq C \|u_{xx}\|_2 \|u_x\|_2^3. \quad (3.12)$$

From (3.7), (3.11), (3.12) and Young's inequality, we have

$$\|u_{xx}(T_1)\|_2 \leq C \quad \text{for any } T_1. \quad \square$$

Now it is possible to present the uniform estimates on smooth solutions for (2.1).

Lemma 3.2. *If u_ε is a smooth solution for (2.1) on $[0, T]$, then there exist a constant $C > 0$ depending only on $\|u_0\|_{H^2}$ and T such that*

$$\|u_\varepsilon(t)\|_{H^2(\Omega)} \leq C, \quad 0 \leq t \leq T.$$

Proof. (1) From Lemma 2.1 we infer that $|u_\varepsilon| = 1$.

(2) As the same procedure as step (2) in Lemma 3.1, we have

$$\begin{aligned} -\frac{d}{dt} \int_{\Omega} |u_{\varepsilon x}|^2 dx &= \varepsilon \int_{\Omega} |u_{\varepsilon xxx}|^2 dx - 3\varepsilon \int_{\Omega} |u_{\varepsilon xx}|^2 u_\varepsilon \cdot u_{\varepsilon xx} dx \\ &\quad - 4\varepsilon \int_{\Omega} (u_{\varepsilon xx} \cdot u_{\varepsilon xxx})(u_\varepsilon \cdot u_{\varepsilon xx}) dx. \end{aligned}$$

So

$$\frac{d}{dt} \|u_{\varepsilon x}\|_2^2 + \varepsilon \|u_{\varepsilon xxx}\|_2^2 \leq 3 \|u_{\varepsilon xx}\|_3^3 + 4\varepsilon \int_{\Omega} |u_{\varepsilon x}| |u_{\varepsilon xx}| |u_{\varepsilon xxx}| dx. \quad (3.13)$$

Applying Lemma 2.2, we obtain

$$\|u_{\varepsilon xx}\|_3 \leq C \|u_{\varepsilon x}\|_2^{\frac{5}{12}} (\|u_{\varepsilon xxx}\|_2 + \|u_{\varepsilon x}\|_2)^{\frac{7}{12}},$$

and

$$\|u_{\varepsilon xx}\|_2 \leq C \|u_{\varepsilon x}\|_2^{\frac{1}{4}} (\|u_{\varepsilon xxx}\|_2 + \|u_{\varepsilon x}\|_2)^{\frac{3}{4}}.$$

Substituting the above inequalities into (3.13), we have

$$\frac{d}{dt} \|u_{\varepsilon x}\|_2^2 + \frac{\varepsilon}{2} \|u_{\varepsilon xxx}\|_2^2 \leq C\varepsilon (1 + \|u_{\varepsilon x}\|_2^{10}).$$

When ε is small enough, we have

$$\|u_{\varepsilon x}(t)\|_2 \leq C, \quad (3.14)$$

where $0 \leq t \leq T$.

(3) As the same procedure as step (3) in Lemma 3.1, we have

$$\begin{aligned} & 4\frac{d}{dt}\|u_{\varepsilon xx}\|_2^2 - 5\frac{d}{dt}\|u_{\varepsilon x}\|_4^4 + 8\varepsilon \int_{\Omega} |u_{\varepsilon xxx}|^2 dx \\ & \leq C\varepsilon \int_{\Omega} |u_{\varepsilon} \cdot u_{\varepsilon xxx}|^2 dx + C\varepsilon \int_{\Omega} |u_{\varepsilon x}|^2 |u_{\varepsilon x} \cdot u_{\varepsilon xxx}| dx \\ & \quad + C\varepsilon \int_{\Omega} |u_{\varepsilon x}|^2 (u_{\varepsilon xx} \cdot u_{\varepsilon})(u_{\varepsilon} \cdot u_{\varepsilon xxx}) dx. \end{aligned}$$

Applying Lemma 2.2, we obtain

$$\|u_{\varepsilon xx}\|_{\infty} \leq C\|u_{\varepsilon x}\|_2^{\frac{1}{2}} \left(\|u_{\varepsilon xxx}\|_2 + \|u_{\varepsilon x}\|_2 \right)^{\frac{1}{2}},$$

and

$$\|u_{\varepsilon x}\|_4 \leq C\|u_{\varepsilon xx}\|_2^{\frac{15}{16}} \left(\|u_{\varepsilon xxx}\|_2 + \|u_{\varepsilon x}\|_2 \right)^{\frac{1}{16}}.$$

By virtue of (3.14), Young's inequality and Hölder's inequality, we have

$$4\frac{d}{dt}\|u_{\varepsilon xx}\|_2^2 - 5\frac{d}{dt}\|u_{\varepsilon x}\|_4^4 \leq C\varepsilon. \quad (3.15)$$

Integrating (3.15) from 0 to t , we have

$$4\|u_{\varepsilon xx}(t)\|_2^2 - 4\|u_{\varepsilon xx}(0)\|_2^2 - 5\|u_{\varepsilon x}(t)\|_4^4 + 5\|u_{\varepsilon x}(0)\|_4^4 \leq C\varepsilon. \quad (3.16)$$

By (3.12), we infer from (3.16) that

$$\|u_{\varepsilon xx}(t)\|_2 \leq C, \quad 0 \leq t \leq T. \quad \square$$

Next we shall deal with the limiting case. Lemma 3.2 and Theorem 5.1 of Chapter 1 [10] imply that there exists a subsequence of u_{ε} (still denoted by u_{ε}) such that when $\varepsilon \rightarrow 0$, there holds

$$u_{\varepsilon} \rightarrow u \quad \text{strongly in } L^p(\Omega \times (0, T)), \quad (3.17)$$

and

$$u_{\varepsilon x} \rightarrow u_x \quad \text{strongly in } L^p(\Omega \times (0, T)), \quad (3.18)$$

where $1 \leq p < \infty$.

Furthermore, we have

$$u_{\varepsilon xx} \rightarrow u_{xx} \quad \text{weakly* in } L^{\infty}(0, T; L^2(\Omega)). \quad (3.19)$$

With the help of these convergence results, taking the scalar product with test function φ in (2.1) and integrating on $\Omega \times (0, T)$, we conclude Theorem 1.1 from the integration by parts.

References

- [1] M. Lakshmanan, K. Porsezian, M. Daniel, Effect of discreteness on the continuum limit of the Heisenberg spin chain, Phys. Lett. A 133 (1988) 483–488.
- [2] K. Porsezian, M. Daniel, M. Lakshmanan, On the integrability aspects of the one-dimensional classical continuum isotropic Heisenberg spin chain, J. Math. Phys. 33 (5) (1992) 1807–1816.

- [3] L.D. Landau, E.M. Lifshitz, On the theory of the dispersion of magnetic permeability in ferromagnetic bodies, *Phys. Z. Sowjetunion* 8 (1935), reproduced in: D. Ter Hear (Ed.), *Collected Papers of L.D. Landau*, Pergamon, New York, 1965, pp. 101–114.
- [4] J. Fivez, On the continuum limit of a classical compressible Heisenberg chain, *Solid State Phys.* 15 (1982) 641–643.
- [5] M. Cieplak, L.A. Turski, Magnetic solitons and elastic kink-like excitations in compressible Heisenberg chain, *J. Phys. C (Solid State)* 13 (1980) L777–L780.
- [6] S. Ding, B. Guo, F. Su, Measure-valued solution to the compressible Heisenberg chain equations, *J. Math. Phys.* 40 (3) (1999) 1153–1162.
- [7] M. Daniel, L. Kavitha, R. Amuda, Soliton spin excitations in an anisotropic Heisenberg ferromagnet with octupole-dipole interaction, *Phys. Rev. B* 59 (21) (1999) 13774–13781.
- [8] Yulin Zhou, Boling Guo, Shaobin Tan, Existence and uniqueness of smooth solution for system of ferromagnetic chain, *Sci. China Ser. A* 34 (1991) 257–266.
- [9] Y.L. Zhou, Interpolation formulas of intermediate quotients for discrete functions with several indices, *J. Comput. Math.* 2 (1984) 276–281.
- [10] J.L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod/Gauthier-Villars, Paris, 1969.